






MISE EN PLACE DU ^{68}Ga -DOTA-Tyr-OCtréotide (edotréotide, Somakit)



Virginie Kouyoumdjian
Denis Mariano-Goulart

$^{68}_{31}\text{Ga}$, un équivalent TEP du $^{99\text{m}}\text{Tc}$

- Cyclotron: $^{69}_{31}\text{Ga} + {}^1_1\text{p} \rightarrow {}^{68}_{32}\text{Ge} + 2. {}^1_0\text{n}$
- $^{68}_{32}\text{Ge} + {}^0_{-1}\text{e} \xrightarrow{270\text{ j}} \nu + {}^{68}_{31}\text{Ga} \xrightarrow{68\text{ min}} {}^{68}_{30}\text{Zn} + {}^0_1\text{e} + \nu$
- Emetteur β^+ (836; 1899 keV) à 88% + CE
et γ (1,08 MeV)
- Ga^{3+} , complexation (DOTA, NOTA,...)




Principaux MRP au ^{68}Ga :

- SSTR: **DOTA-TOC**/NOC/TATE/(AT1S/AT2S) Agoniste SSTR 2 (5/TOC)
- PSMA
- En recherche clinique:
 - Bombésine (Gastrin Releasing Peptide-R)
 - Gliomes, cancer de la prostate
 - α -Melanocyte Stimulating H (mélanome)
 - Cyclo-RGD (Arg-Gly-Asp) (néoangiogénèse tumorale)
 - Antagoniste des R type 4 aux chémokines: (leucémies, sein, prostate, poumons, ovaires, colorectal)
 - Affibody- ^{68}Ga -HER₂⁺ (sein)
 - Biphosphonates
 - Seul: Infection (Semin Nucl Med 2016; 46:436-447)

Enceinte H700 avec automate de synthèse

Société Trasis

Automate de synthèse :
Mini All in One



Iso 5 Classe A avec ventilation et
filtration autonome
Activimètre/ SAS ventilé entrée
sortie matériel

Iso 7 Classe C avec ventilation et
filtration autonome

**Emplacement pour 2
générateurs**: Ger68/Ga68
Place pour bouteille de déchets
liquides dans son blindage

Tiroirs de rangement

Marchepied

Synthèse du ^{68}Ga -DOTA-TOC

- $T(^{68}\text{Ge}) = 270 \text{ j} \Rightarrow A = A_0 \cdot 2^{-t/270}$ en équilibre séculaire
- Rendement d'élution = 60%
- Perte de 4% lors du CQ
- Temps de synthèse + CQ = 50' : $2^{-50/68} = 60\%$
- $A(\text{Ga éluable}) = A_0(\text{Ge}) \cdot 2^{-t/270} \cdot (0,6) \cdot (0,96) \cdot (0,6)$
- Activité par patient : 150 MBq au T1 puis 120 MBq

MOIS	0	1	2	3	4	5	6	7	8	9	10	11	12
A(Ga) éluable (MBq)	629	581	538	497	460	425	393	363	336	311	287	266	246
Nombre de patients	4,2	3,9	3,6	4,1	3,8	3,5	3,3	3,0	2,8	2,6	2,4	2,2	2,0
Nombre de patients	4	4	4	4	4	4	3	3	3	3	3	3	3
Poids moyen à 1,5 MBq/kg	105	97	90	83	77	71	87	81	75	69	64	59	55

Calibration Ge	1850	MBq	
A(Ge) 1° élution	1817	MBq	à 7 jours
REND ELUTION	0,6		
REND CQ	0,96		
TEMPS SYNTH	50	min	
MBq(Ga)/Patient	150	MBq	3 mois
MBq(Ga)/Patient	120	MBq	7 mois

donc changement de générateur
tous les 10 mois

PROTOCOLES

- Laxatif et jeûne non nécessaires
- $\updownarrow$ octréotide discuté (RCP: $\updownarrow$ le jour de l'examen)
- 1,5 (à 3) MBq/kg IVL $\nrightarrow$ Kt nutrition parentérale
- Attente 45' (< 90') post IV puis toilettes
 - Activité maxi à 70 ± 20 min post IV
- Acquisition cTAP de 20' sauf PPGL_{métas/récidive}
- Excrétion rénale
- $E < 3$ mSv
- Suspendre l'allaitement 12h post IV


INTERPRETATION > BDF Foie

FN:

- $\varnothing < 2$.LMH = 8 mm
- dédifférenciée (FDG)

FP :

- lymphocytes activés par inflammation
- ilots (de la tête du) pancréas.
- Adénopathies/K, méningiomes, dysplasie fibreuse.




± Estomac et pancréas

INDICATIONS: Tumeurs > endoderme

- **^{68}Ga -DOTA-TOC**
 - BI **TNE (foregut: bouche→Vater)** $\neq$ carcinome (FDG)
 - TNE thymiques,
 - carcinoïdes bronchiques (^{18}F FDG pour CP petites/grandes cellules)
 - digestives (estomac-D2, Foie, VB, pancréas, rate)
 - Gastrinome, insulinome, glucagonome, VIPome...
 - TNE jéjuno-iléale (midgut) si **^{18}F -DOPA** négative
 - Paragangliomes
 - Phéochromocytomes si **^{18}F -DOPA** négative
 - Méningiomes
- **^{68}Ga -PSMA-11** : $\uparrow$ PSA /K prostate $\pm$ choline normale

NEOPLASIES NEUROENDOCRINES

HORS PPGL: QUEL TRACEUR ?



OCTREOTIDE ET ^{68}Ga -DOTA-TOC

Méta-analyse vs. ^{111}In -OCTREOTIDE
sur 1561 TNE: 44% Δ (30% après In)

^{68}Ga -DOTA-TOC

3,6 fois moins irradiant



Résolution 15 $\rightarrow$ 4 mm

Plus sensible


Acquisition unique de 20'

Pas de laxatif

Pas d'obligation $\updownarrow$ octréotide




^{68}Ga -DOTA-TOC & TNE ILEALE



Rechercher: Médiastin, RP, surrénales + orbites, m. hypoglosse, thyroïde, cœur & péricarde, foie & VB, queue de cheval, paroi vessie (furosémide)...

PHEOCHROMOCYTOME & PARAGANGLIOME: QUEL TRACEUR ?



*si CT/MRI + biochimie (adrénaline) insuffisants

**dont SDHD hérités du père. Parasympathique, T&C + Méd antérieur/moyen

***i.e. périaortique et RP

Table 1 Characteristics of the various hereditary forms of pheochromocytomas and paragangliomas

	First manifestation	Context at presentation (in index cases)	PHEO at presentation	Additional extra-adrenal PGL	Biochemical phenotype	Malignancy risk	Other tumour types
<i>MEN2</i>	MTC	Adult Possible phenotypic features of MEN2 Family history of MTC/PHEO is common	Uni- or bilateral	Very rare	EPI	Very low	Parathyroid adenoma/hyperplasia (MEN2A)
<i>NF1</i>	Neurofibromas	Adult Phenotypic features of neurofibromatosis type 1 Family history of neurofibromatosis type 1 is possible	Often unilateral	Very rare	EPI	Very low	Peripheral nerve sheath tumours, optical glioma ^a
<i>MAX</i>	PHEO	Young adult Family history of PHEO is common	Bilateral	Uncommon	EPI and NE	Moderate	Renal oncocytoma
<i>VHL</i>	PHEO/PGL	Young adult Family history of PPGL is common	Uni- or bilateral	Moderate (retroperitoneum)	NE	Low	Multiple ^b
<i>SDHA</i>	PHEO/PGL	Adult Family history of PPGL is extremely rare	Rare	Moderate (retroperitoneum or head and neck)	NE and/or DA	Moderate/high	GIST ^c , RCC
<i>SDHB</i>	PHEO/PGL	Adult Family history of PPGL is possible	Often unilateral	Common (retroperitoneum)	NE and/or DA	High	GIST ^c , pituitary adenoma, RCC
<i>SDHD</i>	PHEO/PGL	Adult Family history of HNPGGL is common	Uni- or bilateral	Common (head and neck)	NE and/or DA	Moderate	GIST ^c , pituitary adenoma, RCC
<i>SDHC</i>	PHEO/PGL	Adult Family history of PHEO/PGL is rare	Rare	Common (mediastinum)	NE	Low/Moderate	GIST ^c , RCC
<i>HIF2A</i>	Polycythaemia (often at birth or in early childhood)	Adolescent girls or young adult women Family history of PPGL is absent	Rare	Expected (retroperitoneum)	NE	High	Somatostatinoma


MAX: myc-associated factor X, MEN: multiple endocrine neoplasia, NF1: neurofibromin 1, VHL: von Hippel–Lindau, SDHA, -B, -C, and -D: succinate dehydrogenase subunits A, B, C, and D, HIF2A: hypoxia-inducible factor 2α, PPGL: pheochromocytoma and paraganglioma, PHEO: pheochromocytoma, PGL: paraganglioma, MTC: medullary thyroid carcinoma, HNPGGL: head and neck paraganglioma; GIST: gastrointestinal stromal tumour, RCC: renal cell carcinoma, EPI: epinephrine, NE: norepinephrine, DA: dopamine

^a *NF1*-associated PGEs/PGLs are characterised by the presence of multiple neurofibromas, café-au-lait spots, Lisch nodules of the iris, and other rare disorders



^b *VHL* disease is an autosomal dominant disorder that also predisposes to renal tumours and clear cell carcinoma, pancreatic serous cystadenomas, pancreatic neuroendocrine tumours, testicular tumours, and haemangioblastoma of the eye and central nervous system

^c Non-*KIT/PDGFRA* GISTs may be caused by mutations in the *SDHA-D* genes and could be associated with PGLs in patients with Carney–Stratakis syndrome

^d *HIF2A*-related PHEOs/PGLs are associated with the presence of duodenal somatostatinoma and retinal abnormalities (Pacak–Zhuang syndrome)



EXEMPLES



[⁶⁸Ga]-DOTATATE

[¹⁸F]-FDG PET

[¹⁸F]-FDOPA



PGL métastatique (poumon, os) SDHB +

^{18}F -CHOLINE ET ^{68}Ga -PSMA





^{68}Ga -PSMA-11: ligand inhibiteur du PSMA, protéine transmembranaire $\uparrow$ ADC prostate $\propto$ Gleason, gravité
Pas si spé: grêle, pancréas, SNP, glandes salivaires...

ATU 2018: $\uparrow$ PSA $\pm$ choline normale
10 % d'examens TEP-Ga-PSMA normaux
30-50% changement thérapeutique






^{68}Ga -PSMA après ^{18}F -CHOLINE négative

FDG & CANCERS DEDIFFERENCIES




Se (^{99m}Tc -BP) >> Se(^{18}F -DG)
Sauf si méta agressive, dédifférenciée



Neuroblastome

MERCI POUR VOTRE ATTENTION



v-kouyoumdjian@chu-montpellier.fr
d-mariano_goulart@chu-montpellier.fr